

Bruck decomposition for endomorphisms of quasigroups

Péter T. Nagy (Debrecen) and Peter Plaumann (Cuernavaca)

April 30, 2009

Abstract

In the year 1944 R. H. Bruck has described a very general construction method which he called the extension of a set by a quasigroup. We use it to construct a class of examples for LF -quasigroups in which the image of the map $e(x) = x \setminus x$ is a group. More generally, we consider the variety of quasigroups which is defined by the property that the map e is an endomorphism and its subvariety where the image of the map e is a group. We characterize quasigroups belonging to these varieties using their Bruck decomposition with respect to the map e .

1 Introduction

A binary algebra $(Q, \cdot)$ with multiplication $(x, y) \mapsto x \cdot y$ is called a *quasigroup* if the equations $a \cdot y = b$ and $x \cdot a = b$ have precisely one solution in Q which we denote by $y = a \setminus b$ and $x = b / a$. The elements $1_l(a) = a / a$ respectively $1_r(a) = a \setminus a$ are the left respectively the right local unit element of the element a . If the left (right) local unit elements coincide for all elements of $(Q, \cdot)$, then the element $1_l = 1_l(a)$, resp. $1_r = 1_r(a)$, is called the left, resp. right, unit element of $(Q, \cdot)$. If a quasigroup $(Q, \cdot)$ has both left and right unit elements then they coincide $1 = 1_l = 1_r$, in this case $(Q, \cdot)$ is called a *loop*.

In the year 1944 R. H. Bruck has described a very general construction method which he called the extension of a set by a quasigroup (cf. [3], [4]). Epimorphisms of quasigroups in general cannot be described by cosets of a normal subquasigroup, but only by congruence relations in the sense of universal algebra. Bruck's construction takes this into account giving a manageable description of quasigroup epimorphisms. In this note we discuss

2000 *Mathematics Subject Classification*: 20N05 .

Key words and phrases: Bruck extension of quasigroups, non-associative extension of groups, special varieties of quasigroups, LF -quasigroups

This research was supported by the Hungarian Scientific Research Fund (OTKA), Grant K 67617 and by the Program *Cátedras Especiales. SRE* of the Mexican Government.

this method for endomorphisms of quasigroups.

A quasigroup $(Q, \cdot)$ is called an *LF-quasigroup* if the identity

$$x \cdot yz = xy \cdot (x \backslash x \cdot z)$$

holds in Q . In his book [1] V. D. Belousov initiated a systematic study of *LF-quasigroups* using isotopisms. Recently progress has been made in this topic (cf. [9], [7], [8]). It is known that in an *LF-quasigroup* the map $e(x) = x \backslash x$ is an endomorphism, which we call the left deviation. In this situation Bruck's theory is available. We use it to construct a class of examples for *LF-quasigroups* Q in which $e(Q)$ is a group.

More generally, we consider the variety $\mathfrak{D}_l$ of quasigroups which is defined by the property that the left deviation is an endomorphism and its subvariety $\mathfrak{a}D_l$ where the image of the left deviation is a group. We characterize quasigroups belonging to these varieties using their Bruck decomposition with respect to their left deviation.

2 The Bruck decomposition of a quasigroup

In Bruck's papers [3], (Theorem 10 A, pp. 166-168) and [4], (pp. 778-779) a principal construction for quasigroups is given. Let $(E, \cdot, \backslash, /)$ be a quasigroup, let T be a set and let $\{\nabla_{a,b}; a, b \in E\}$ be a family of multiplications on T . Define on the set $Q = T \times E$ a multiplication by

$$(\alpha, a) \circ (\beta, b) = (\alpha \nabla_{a,b} \beta, ab), \quad (\alpha, a), (\beta, b) \in T \times E. \quad (1)$$

Then $(Q, \circ)$ is a quasigroup if and only if for any $a, b \in E$ the multiplication $\nabla_{a,b}$ on T defines a quasigroup $\mathcal{T}_{a,b} = (T, \nabla_{a,b})$. In this case we call $\mathcal{B} = (E, T, (\nabla_{a,b})_{a,b \in E})$ *Bruck system* and put $Q(\mathcal{B}) = Q$. Obviously, the projection $((\alpha, a) \mapsto a) : Q(\mathcal{B}) \rightarrow E$ is an epimorphism of quasigroups. (Cf. [6], pp. 35-36.) We call this epimorphism the *canonical epimorphism* for $\mathcal{B}$.

Conversely, let $(Q, \circ)$ and $(E, \cdot)$ be quasigroups and let $\pi : Q \rightarrow E$ be an epimorphism. For the inverse images $T_a = \pi^{\leftarrow}(a)$, $a \in E$, one has

$$T_a \circ T_b = T_{ab},$$

hence the set $\{T_a, a \in E\}$ forms a quasigroup $\mathcal{E}'$ isomorphic to $(E, \cdot)$. Using a transversal for the partition $\{T_a, a \in E\}$ of the set Q one can identify the inverse images T_a , $a \in E$, with a subset $T \subset Q$ and the set Q with the cartesian product $T \times E$. The multiplication in Q can be written in the form (1), where $\alpha, \beta \in T$, $a, b \in E$. This means that we have obtained a Bruck system $\mathcal{B}_\pi = (E, T, (\nabla_{a,b})_{a,b \in E})$ for which $Q(\mathcal{B}_\pi)$ is isomorphic to Q and which has π as canonical epimorphism. We call this representation of $(Q, \circ)$ a *Bruck decomposition of $(Q, \circ)$ with respect to $\pi : Q \rightarrow E$* .

Let $\eta : Q \rightarrow Q$ be an endomorphism. We consider the set E of the congruence classes $T_x = \eta^\leftarrow(x)$, $x \in \eta(Q)$. The multiplication $T_x \star T_y = T_{xy}$ defines a quasigroup $(E, \star)$ such that the mapping $\pi : (x \mapsto T_x) : Q \rightarrow E$ is an epimorphism. We put $\iota = \eta \circ \pi^\leftarrow = (T_x \mapsto \eta(x)) : E \rightarrow Q$. Then ι is an injective homomorphism (see Theorem 6.12, [5], p. 50).

We consider the Bruck decomposition $Q = T \times E$ with respect to the epimorphism π . For $a \in E = \{T_x, x \in \eta(Q)\}$ one has $\iota(a) = (\gamma(a), g(a))$, where $\gamma(a) \in T$, $g(a) \in E$. The maps $\gamma : E \rightarrow T$ and $g : E \rightarrow E$ satisfy

$$\begin{aligned} (\gamma(ab), g(ab)) &= \iota(ab) = \iota(a)\iota(b) = \\ &= (\gamma(a), g(a))(\gamma(b), g(b)) = (\gamma(a) \nabla_{g(a), g(b)} \gamma(b), g(a)g(b)). \end{aligned}$$

It follows that $g : E \rightarrow E$ is an endomorphism and

$$\gamma(ab) = \gamma(a) \nabla_{g(a), g(b)} \gamma(b) \quad (2)$$

holds for all $a, b \in E$. We call the structure described here the *Bruck decomposition of the quasigroup $(Q, \circ)$ with respect to the endomorphism $\eta : Q \rightarrow Q$* .

If the quasigroup Q is a loop and $\eta : Q \rightarrow Q$ is an endomorphism, then $K = \eta^\leftarrow(1)$ is a normal subloop of Q . In this situation Q is a semidirect product of K and $\eta(Q)$ if and only if $\eta = \eta^2$ holds. We describe the Bruck decomposition with respect to an idempotent endomorphism for arbitrary quasigroups.

Proposition 1 *An endomorphism η of a quasigroup Q is idempotent if and only if in the Bruck decomposition with respect to η the maps $\gamma : E \rightarrow T$ and $g : E \rightarrow E$ satisfy*

$$g^2 = g \quad \text{and} \quad \gamma \circ g = \gamma,$$

i. e. if and only if the endomorphism $g : E \rightarrow E$ is idempotent and the map $\gamma : E \rightarrow T$ factors over the congruence relation defined by g on E .

Proof. Since $\eta(\alpha, a) = \iota(\pi(\alpha, a)) = \iota(a) = (\gamma(a), g(a))$ the assertion follows from

$$\eta(\eta(\alpha, a)) = \eta(\gamma(a), g(a)) = (\gamma(g(a), g(g(a)))).$$

■

3 The left deviation of a quasigroup

For a quasigroup $(Q, \cdot, \backslash, /)$ we call the map $e = (x \mapsto x \backslash x) : Q \rightarrow Q$ the *left deviation*. As mentioned in the preliminaries the left deviation of $x \in Q$ is

the local right unit element of x . In a Bruck decomposition $Q = T \times E$ with respect to an epimorphism $Q \rightarrow E$ the deviation is $e(\alpha, a) = (\alpha \backslash \alpha, a \backslash a)$, where $a \backslash a$ is computed in the quasigroup E and $\alpha \backslash \alpha$ is computed in the quasigroup $\mathcal{T}_{a, a \backslash a} = (T, \nabla_{a, a \backslash a})$.

Obviously, the quasigroups in which *the left deviation is an endomorphism form a variety* $\mathfrak{D}_l$ of quasigroups. For $(Q, \cdot, \backslash, /) \in \mathfrak{D}_l$ we consider the Bruck decomposition $Q = T \times E$ with respect to the left deviation. In this case $e(\alpha, a) = (\gamma(a), g(a))$ and hence $g(a) = a \backslash a$ (computed in E) and

$$\alpha \nabla_{a, g(a)} \gamma(a) = \alpha. \quad (3)$$

Theorem 2 *A quasigroup Q belongs to the variety $\mathfrak{D}_l$ if and only if there exists a Bruck system $\mathcal{B} = (E, T, (\nabla_{a, b})_{a, b \in E})$ satisfying*

- (i) $Q \cong Q(\mathcal{B})$,
- (ii) *the quasigroup $\mathcal{T}_{a, a \backslash a} = (T, \nabla_{a, a \backslash a})$ has a right unit element, denoted by $\epsilon(a)$, for any $a \in E$,*
- (iii) *the map $(a \mapsto (\epsilon(a), a \backslash a)) : E \rightarrow Q(\mathcal{B})$ is a homomorphism.*

In this case the left deviation of $Q(\mathcal{B})$ is the map

$$e = ((\alpha, a) \mapsto (\epsilon(a), a \backslash a)) : Q(\mathcal{B}) \rightarrow Q(\mathcal{B}).$$

Proof. Assume first that Q belongs to $\mathfrak{D}_l$ and consider the Bruck decomposition with respect to the left deviation $e(x) = x \backslash x$ of Q . Then E is isomorphic to the subquasigroup $e(Q)$ of Q . Putting $\epsilon = \gamma$ the assertion (ii) follows from equation (3) and the assertion (iii) follows from equation (2). Conversely, if $\mathcal{B} = (E, T, (\nabla_{a, b})_{a, b \in E})$ is a Bruck system satisfying (i) then $e(\alpha, a) = (\alpha, a) \backslash (\alpha, a) = (\alpha', a \backslash a)$, where $\alpha = \alpha \nabla_{a, a \backslash a} \alpha'$. From (ii) it follows that $\alpha' = \epsilon(a)$ and the deviation satisfies $e(\alpha, a) = (\epsilon(a), a \backslash a)$. Hence we obtain from (iii) that the quasigroup Q belongs to the variety $\mathfrak{D}_l$. ■

Corollary 3 *Let $\mathcal{B} = (E, T, (\nabla_{a, b})_{a, b \in E})$ be a Bruck system satisfying the conditions (ii) and (iii) of the previous theorem. Then $\mathcal{B}$ is a Bruck decomposition of the quasigroup $Q(\mathcal{B})$ with respect to the left deviation of $Q(\mathcal{B})$ if and only if the homomorphism*

$$\iota = (a \mapsto (\epsilon(a), a \backslash a)) : E \rightarrow Q(\mathcal{B})$$

is injective.

Example 4 Let $(E, \cdot)$, $\mathcal{T}^{(1)} = (T, \circ)$ and $\mathcal{T}^{(2)} = (T, \star)$ be quasigroups such that the following properties are satisfied:

- (i) E is a $\mathfrak{D}_l$ -quasigroup,
- (ii) $\mathcal{T}^{(1)}$ has a right unit element ϵ ,
- (iii) ϵ is idempotent in the quasigroup $\mathcal{T}^{(2)}$.

Put

- (i) $\mathcal{T}_{a,a \setminus a} = (T, \nabla_{a,a \setminus a}) = \mathcal{T}^{(1)} = (T, \circ)$,
- (ii) $\mathcal{T}_{a \setminus a, b \setminus b} = (T, \nabla_{a \setminus a, b \setminus b}) = \mathcal{T}^{(2)} = (T, \star)$ if $b \setminus b \neq (a \setminus a) \setminus (a \setminus a)$,
- (iii) $\mathcal{T}_{c,d} = (T, \nabla_{c,b \setminus d})$ arbitrary in all other cases.

According to Theorem 2 the multiplication $(\alpha, a) \circ (\beta, b) = (\alpha \nabla_{a,b} \beta, ab)$ on the set $T \times E$ is a $\mathfrak{D}_l$ -quasigroup Q with left deviation $e(\alpha, a) = (\epsilon, a \setminus a)$. For $(E, \cdot)$ one can take idempotent quasigroups or groups. Clearly, the construction of the quasigroup $Q = T \times E$ gives a Bruck decomposition with respect to the left deviation of Q if and only if the left deviation of $(E, \cdot)$ is an automorphism. If $(E, \cdot)$ is a group with unit element 1 then the left deviation of Q is the constant $(\epsilon, 1)$ which is the right unit element of Q .

4 Associative image of the left deviation map

The class of $\mathfrak{D}_l$ -quasigroups for which the image of the left deviation map is a group forms a variety, too, as can be seen from the identities:

$$xy \setminus xy = x \setminus x \cdot y \setminus y, \quad (x \setminus x \cdot y \setminus y) \cdot z \setminus z = x \setminus x \cdot (y \setminus y \cdot z \setminus z). \quad (4)$$

We denote this variety by $\mathfrak{aD}_l$ and investigate their Bruck decomposition. As an immediate consequence of Theorem 2 one obtains

Theorem 5 *A quasigroup Q belongs to the variety $\mathfrak{aD}_l$ if and only if there exists a Bruck system $\mathcal{B} = (E, T, (\nabla_{a,b})_{a,b \in E})$ with $Q \cong Q(\mathcal{B})$ satisfying*

- (i) E is a group (with unit element 1),
- (ii) the quasigroup $\mathcal{T}_{a,1} = (T, \nabla_{a,1})$ has right unit element, denoted by $\epsilon(a)$, for any $a \in E$,
- (iii) the map $\epsilon : E \rightarrow \pi^{\leftarrow}(1) = \mathcal{T}_{1,1} = (T, \nabla_{1,1})$ is a homomorphism of the group E into the normal subquasigroup $\pi^{\leftarrow}(1) = \mathcal{T}_{1,1} = (T, \nabla_{1,1})$ of $Q(\mathcal{B})$.

In this case one has $e(\alpha, a) = (\epsilon(a), 1)$ for any $(\alpha, a) \in T \times E$. ■

Proof. E is a group since it is isomorphic to the image of the left deviation map. Hence $e(\alpha, a) = (\epsilon(a), a \backslash a) = (\epsilon(a), 1)$ for any $(\alpha, a) \in T \times E$ and $\epsilon : (a \mapsto (\epsilon(a), 1)) : E \rightarrow \pi^{\leftarrow}(1) = \mathcal{T}_{1,1}$. ■

Example 6 Let $(E, \cdot), (T, \circ)$ be quasigroups and let $\epsilon : E \rightarrow T$ be a homomorphism such that the following properties are satisfied:

- (i) $(E, \cdot)$ is a group with unit element 1,
- (ii) $\epsilon(1)$ is a right unit element of $(T, \circ)$.

Put

- (a) $\alpha \nabla_{a,1} \beta = (\alpha / \epsilon(a)) \circ \beta$ in the quasigroup $\mathcal{T}_{a,1} = (T, \nabla_{a,1})$, where $/$ is the right division in $(T, \circ)$,
- (b) $\mathcal{T}_{a,b} = (T, \nabla_{a,b})$ arbitrary for $b \neq 1$.

According to Theorem 5 the multiplication $(\alpha, a) \circ (\beta, b) = (\alpha \nabla_{a,b} \beta, ab)$ on the set $T \times E$ is an $\mathbf{aD}_l$ -quasigroup. For $(E, \cdot)$ and $(T, \circ)$ one can take groups having $\epsilon : E \rightarrow T$ as a group homomorphism. The decomposition $Q = T \times E$ is a Bruck decomposition with respect to the left deviation of Q if and only if the homomorphism $\epsilon : E \rightarrow T$ is injective.

5 LF -quasigroups

It is known that the LF -quasigroups form a subvariety of $\mathfrak{D}_l$ (cf. [2] p. 108. and Lemma 2.1. in [9]). We shall now give examples of LF -quasigroups even belonging to the variety $\mathbf{aD}_l$.

Example 7 Let E, T be groups and let $\epsilon : E \rightarrow T$ be a homomorphism. Put

$$\alpha \nabla_{a,b} \beta = \alpha \cdot \epsilon(a)^{-1} \cdot \beta$$

for all $a, b \in E$. Then every $\mathcal{T}_{a,b} = (T, \nabla_{a,b})$ is a group (with unit element $\epsilon(a)$) which is isotopic and hence isomorphic to the group T .

As in the previous example the multiplication $(\alpha, a) \circ (\beta, b) = (\alpha \nabla_{a,b} \beta, ab)$ on the set $T \times E$ defines an $\mathbf{aD}_l$ -quasigroup (with left unit element $(1, 1)$) in which the left deviation is given by $e(\alpha, a) = (\epsilon(a), 1)$ (Theorem 5).

Theorem 8 Let E, T be groups and let $\epsilon : E \rightarrow T$ be a homomorphism. The set $Q = T \times E$ equipped with multiplication $(\alpha, a) \circ (\beta, b) = (\alpha \cdot \epsilon(a)^{-1} \cdot \beta, ab)$ is an LF -quasigroup with left unit element $(1, 1)$ satisfying the left inverse property.

Proof. An easy calculation shows

$$(\alpha, a) \circ ((\beta, b) \circ (\gamma, c)) = (\alpha \cdot \epsilon(a)^{-1} \cdot \beta \cdot \epsilon(b)^{-1} \cdot \gamma, abc).$$

On the other hand

$$\begin{aligned} ((\alpha, a) \circ (\beta, b)) \circ ((\epsilon(a), 1) \circ (\gamma, c)) &= (\alpha \cdot \epsilon(a)^{-1} \cdot \beta, ab) \circ (\epsilon(a) \cdot \gamma, c) = \\ &= (\alpha \cdot \epsilon(a)^{-1} \cdot \beta \cdot \epsilon(ab)^{-1} \cdot \epsilon(a) \cdot \gamma, abc). \end{aligned}$$

Hence $(Q, \circ)$ is an LF -quasigroup in which $(1, 1)$ is the left unit element. The left inverse of an element (α, a) is given by $(\epsilon(a) \cdot \alpha^{-1} \cdot \epsilon(a)^{-1}, a^{-1})$. Indeed one has $(\epsilon(a) \cdot \alpha^{-1} \cdot \epsilon(a)^{-1}, a^{-1}) \circ ((\alpha, a) \circ (\beta, b)) = (\beta, b)$. ■

The decomposition $Q = T \times E$ is a Bruck decomposition with respect to the left deviation of Q if and only if the homomorphism $\epsilon : E \rightarrow T$ is injective.

We note that in accordance with Theorem 4.1 in [9] the quasigroup $Q = T \times E$ is isotopic to the direct product of the groups T and E , the isotopism is given by the triple $(\phi, \text{id}, \text{id})$, where $\phi : (\alpha, a) \mapsto (\alpha \cdot \epsilon(a)^{-1}, a)$.

References

- [1] V. D. Belousov, *Foundations of the Theory of Quasigroups and Loops*, Nauka, Moscow, 1967 (in Russian).
- [2] V. D. Belousov, *Elements of Quasigroup Theory: A Special Course*, Kishinev State University Press, Kishinev, 1981.
- [3] R. H. Bruck, *Some Results in the Theory of Linear Non-Associative Algebras*, Trans. Amer. Math. Soc., **56**, (1944), pp. 141-199.
- [4] R. H. Bruck, *Simple Quasigroups*, Bull. Amer. Math. Soc., **50**, (1944), pp. 769-781.
- [5] S. Burris, H. P. Sankappanavar, *A Course in Universal Algebra*, Graduate Texts in Mathematics, Vol. 78. New York - Heidelberg - Berlin, Springer-Verlag, 1981.
<http://www.math.uwaterloo.ca/~snburris/htdocs/UALG/univ-algebra.pdf>
- [6] O. Chein, *Examples and methods of construction*, Chapter II. in Quasigroups and Loops: Theory and Applications, (Eds.: O. Chein – H. O. Pflugfelder – J. D. H. Smith,) pp. 27–93, Heldermann Verlag, Berlin, (1990).
- [7] T. Kepka – M. K. Kinyon – J.D. Phillips, *The structure of F-quasigroups*, Journal of Algebra **317** (2007) pp. 435-461.

- [8] P. Plaumann – L. Sabinina – L. Sbitneva, *A decomposition of LF-quasigroups*, AMS Contemporary Mathematics (to appear).
- [9] V.A. Shcherbacov, *On the structure of left and right F-, SM- and E-quasigroups*, arXiv:0811.1725.

Péter T. Nagy
 Institute of Mathematics
 University of Debrecen
 P.O.B. 12
 4010 DEBRECEN, HUNGARY
E-mail: nagypeti@math.unideb.hu

Peter Plaumann
 Mathematisches Institut der
 Universität Erlangen-Nürnberg
 Bismarckstr. 1 $\frac{1}{2}$
 91054 ERLANGEN, GERMANY
E-mail: peter.plaumann@mi.uni-erlangen.de